


The Bahamas

C. McSweeney¹, M. New^{1,2} and G. Lizcano¹

1. School of Geography and Environment, University of Oxford.
2. Tyndall Centre for Climate Change Research

<http://country-profiles.geog.ox.ac.uk>



General Climate

The Bahamas are an archipelago made up of several hundred islands which are located North of Cuba and the Caribbean Islands, at a latitude of 21-27°N, on the boundary of the tropical and sub-tropical zones. The Islands experience the year-round warm, humid conditions associated with the Tropics, but with more distinct seasonal variations than the Southern Caribbean Islands.

Seasonal mean temperatures range from 21-24°C in the cooler months of December to February, to 27-28°C in the warmer months of July and August. The wet season occurs through May to October, during which the islands receive around 100-150mm per month.

Inter-annual variability in climate here is influenced strongly by the El Niño Southern Oscillation (ENSO). El Niño episodes bring warmer and drier than average conditions between June and August and La Niña episodes bring colder and wetter conditions at this time. The Bahamas also lie in the heart of the Atlantic hurricane belt, where Hurricanes occur throughout August, September and October. Heavy rainfall associated with cyclones and hurricanes contributes significantly to wet season rainfall totals. The occurrence of hurricanes is strongly linked to ENSO, with more frequent hurricane activity associated with La Niña events, and less frequent events in El Niño years.

Recent Climate Trends

Temperature

- Mean annual temperature has increased by around 0.5°C since 1960, at an average rate of 0.11 °C per decade. The rate of increase is most rapid in the warmest seasons (JJA and SON) at 0.13 and 0.15°C per decade respectively. The warming is also more rapid over the north-eastern islands than the south-west.
- There is insufficient daily observational data to identify trends in daily temperature extremes.

Precipitation

- Mean rainfall over The Bahamas has not changed significantly since 1960.

GCM Projections of Future Climate

Temperature

- The mean annual temperature is projected to increase by 0.8 to 2.3°C by the 2060s, and 1.2 to 3.5 degrees by the 2090s. The range of projections by the 2090s under any one emissions scenario is around 1-2°C.
- The projected rate of warming is most rapid in summer (JJA).
- All projections indicate substantial increases in the frequency of days and nights that are considered 'hot'¹ in current climate.
 - Annually, projections indicate that 'hot' days will occur on 24-47% of days by the 2060s, and 26-67% of days by the 2090s. Days considered 'hot' by current climate standards for their season are projected to increase most rapidly in JJA, occurring on 49-99% of days of the season by the 2090s.
 - Nights that are considered 'hot' for the annual climate of 1970-99 are projected to occur on 28-47% of nights by the 2060s and 31-66% of nights by the 2090s. Nights that are hot for each season are projected to increase most rapidly in JJA, occurring on 61-99% of nights in every season by the 2090s.
- All projections indicate decreases in the frequency of days and nights that are considered 'cold'² in current climate. These events are expected to become exceedingly rare, occurring on 0-4% of days in the year, and potentially not at all under the higher emissions scenarios.

Precipitation

- Projections from different models in the ensemble are broadly consistent in indicating decreases in rainfall for The Bahamas. This is mainly due to decreases in MAM and JJA rainfall. Projected changes for these seasons by the 2090s range from -42 to +14% in MAM and -59 to +12% in JJA.
- The proportion of total rainfall that falls in heavy³ events decreases in most projections, for MAM and JJA rainfall, changing by -19 to +7%.
- Maximum 1- and 5-day rainfalls in projections do not show a consistent direction of change, but tend towards decreases in MAM and JJA.

¹ 'Hot' day or 'hot' night is defined by the temperature exceeded on 10% of days or nights in current climate of that region and season.

² 'Cold' days or 'cold' nights are defined as the temperature below which 10% of days or nights are recorded in current climate of that region or season.

³ A 'Heavy' event is defined as a daily rainfall total which exceeds the threshold that is exceeded on 5% of rainy days in current the climate of that region and season.

Additional Regional Climate Change Information

- Tropical cyclones are poorly captured by GCMs and thus potential changes in intensity and tracks of tropical cyclones in the future are very uncertain. Whilst evidence indicates that tropical cyclones are likely to become, on the whole, more intense under a warmer climate as a result of higher sea-surface temperatures, there is great uncertainty in changes in frequency, and changes to storm tracks and their interactions with other features of climate variability (such as the El Niño Southern Oscillation) which introduces uncertainty at the regional scale (Christensen *et al.*, 2007).
- Uncertainty in potential changes in tropical cyclone contributes to uncertainties in future wet-season rainfall. Potential increases in summer rainfall associated with tropical cyclone activity, which may not be captured in the GCM projections, may counteract the projected decreases in rainfall in the region (Christensen *et al.*, 2007).
- Model simulations show wide disagreements in projected changes in the amplitude of future El Niño events, contributing to uncertainty in future climate variability in projections for this region.
- The Bahamas are vulnerable to sea-level rise. Sea-level in this region is projected by climate models to rise by the following levels⁴ by the 2090s, relative to 1980-1999 sea-level:
 - 0.13 to 0.43m under SRES B1
 - 0.16 to 0.53m under SRES A1B
 - 0.18 to 0.56m under SRES A2
- For further information see Christensen *et al.* (2007) IPCC Working Group I Report: '*The Physical Science Basis*', Chapter 11 (*Regional Climate Projections*): Sections 11.6 (*South and Central America*) and 11.9 (*Small Islands*).

⁴ Taken from the IPCC Working group I (*The Physical Science Basis*): Chapter 10 (Global Climate Projections) (Meehl *et al.*, 2007). Regional sea-level projections are estimated by applying regional adjustments (Fig 10.32, p813) to projected global mean sea-level rise from 14 AR4 models.

Data Summary

	Observed Mean 1970-99	Observed Trend 1960-2006	Projected changes by the 2030s			Projected changes by the 2060s			Projected changes by the 2090s				
			Min	Median	Max	Min	Median	Max	Min	Median	Max		
Temperature													
(change in °C per decade)													
<i>Annual</i>													
Annual	25.3	0.11*	A2	0.6	0.9	1.2	1.5	1.8	2.0	2.3	3.1	3.5	
			A1B	0.4	1.0	1.3	1.1	1.8	2.3	1.9	2.5	3.0	
			B1	0.3	0.8	1.0	0.8	1.3	1.6	1.2	1.5	2.2	
DJF	22.6	0.12	A2	0.4	0.9	1.2	1.3	1.6	2.1	2.2	2.8	3.5	
			A1B	0.1	0.9	1.4	1.2	1.7	2.1	1.5	2.2	3.2	
			B1	0.4	0.8	1.1	0.8	1.3	1.9	1.0	1.5	2.1	
MAM	24.3	0.06	A2	0.5	0.9	1.4	1.3	1.7	2.1	2.1	2.9	3.3	
			A1B	0.4	1.0	1.3	1.0	1.8	2.1	1.8	2.5	2.9	
			B1	0.2	0.8	1.1	0.9	1.1	1.4	0.9	1.6	2.1	
JJA	27.9	0.13*	A2	0.7	1.0	1.3	1.6	1.9	2.4	1.9	2.6	3.0	
			A1B	0.5	1.0	1.5	1.1	1.9	2.4	1.2	1.6	2.2	
			B1	0.2	0.8	1.2	0.7	1.4	1.8	2.3	3.0	3.9	
SON	26.5	0.15*	A2	0.8	1.1	1.1	1.4	2.0	2.2	1.9	2.5	3.4	
			A1B	0.5	1.1	1.5	1.0	1.9	2.5	1.3	2.5	2.3	
			B1	0.3	0.8	1.0	0.7	1.4	1.7	1.3	1.6	2.3	
Precipitation													
(mm per month)													
(change in mm per decade)													
<i>Annual</i>													
Annual	99.3	0.6	A2	-14	0	12	-26	-4	12	-32	-7	22	
			A1B	-19	-4	4	-31	-2	16	-18	-6	18	
			B1	-8	-2	18	-9	-1	13	-12	-1	14	
DJF	49.4	-0.9	A2	-16	-1	9	-17	-4	19	-21	0	26	
			A1B	-17	-3	22	-21	0	16	-18	0	32	
			B1	-5	0	13	-14	-1	9	-7	-2	31	
MAM	84.4	3.1	A2	-15	-3	21	-29	-9	9	-31	-14	3	
			A1B	-20	-2	3	-25	-7	9	-26	-10	3	
			B1	-17	-4	17	-17	-8	12	-19	-5	11	
JJA	142.1	0.7	A2	-33	0	7	-42	-12	1	-64	-16	13	
			A1B	-31	-9	0	-49	-12	11	-48	-15	9	
			B1	-17	-7	8	-25	-4	18	-21	-9	14	
SON	121.5	-0.2	A2	-7	1	19	-18	5	43	-35	9	58	
			A1B	-14	1	21	-36	13	34	-15	4	33	
			B1	-17	4	50	-7	9	33	-9	5	28	
Precipitation (%)													
(mm per month)													
(change in % per decade)													
<i>Annual</i>													
Annual	99.3	0.6	A2	-13	-1	10	-25	-4	10	-30	-7	18	
			A1B	-17	-4	3	-29	-2	13	-18	-8	15	
			B1	-8	-3	15	-14	-1	11	-12	-2	12	
DJF	49.4	-1.8	A2	-17	-2	11	-20	-6	22	-20	0	30	
			A1B	-16	-3	18	-22	-1	18	-18	-1	33	
			B1	-6	0	17	-19	-1	11	-6	-3	25	
MAM	84.4	3.7	A2	-27	-8	35	-40	-15	15	-42	-20	5	
			A1B	-28	-5	5	-39	-10	15	-41	-16	5	
			B1	-22	-8	31	-27	-15	22	-31	-6	14	
JJA	142.1	0.5	A2	-30	0	7	-38	-12	1	-59	-22	12	
			A1B	-28	-12	0	-45	-12	10	-44	-17	9	
			B1	-21	-8	8	-32	-4	17	-22	-12	9	
SON	121.5	-0.2	A2	-4	1	10	-11	3	24	-22	5	31	
			A1B	-9	0	14	-23	10	27	-12	2	22	
			B1	-10	2	27	-7	5	26	-6	3	20	

		Observed Mean 1970-99	Observed Trend 1960-2006	Projected changes by the 2030s			Projected changes by the 2060s			Projected changes by the 2090s		
	% Frequency			Min	Median	Max	Min	Median	Max	Min	Median	Max
Frequency of Hot Days (TX90p)						<i>Future % frequency</i>						<i>Future % frequency</i>
Annual	****	****	A2	****	****	****	26	40	47	36	59	67
			A1B	****	****	****	28	39	45	32	49	61
			B1	****	****	****	24	33	38	26	38	45
			A2	****	****	****	37	50	73	61	77	96
DJF	****	****	A1B	****	****	****	41	51	75	46	69	89
			B1	****	****	****	30	38	70	32	50	81
			A2	****	****	****	26	48	67	47	75	95
MAM	****	****	A1B	****	****	****	31	47	69	46	67	92
			B1	****	****	****	22	36	58	22	47	78
			A2	****	****	****	57	73	85	78	96	99
JJA	****	****	A1B	****	****	****	56	73	89	69	93	98
			B1	****	****	****	48	59	69	49	76	84
			A2	****	****	****	33	65	82	53	88	97
SON	****	****	A1B	****	****	****	36	69	81	41	85	93
			B1	****	****	****	27	55	72	35	67	79
Frequency of Hot Nights (TN90p)						<i>Future % frequency</i>						<i>Future % frequency</i>
Annual	****	****	A2	****	****	****	32	40	47	44	57	66
			A1B	****	****	****	33	39	44	39	49	60
			B1	****	****	****	28	33	38	31	38	45
			A2	****	****	****	32	43	64	54	71	92
DJF	****	****	A1B	****	****	****	37	46	67	44	61	83
			B1	****	****	****	29	33	62	30	45	74
			A2	****	****	****	32	43	65	58	71	95
MAM	****	****	A1B	****	****	****	29	45	68	54	63	93
			B1	****	****	****	27	34	56	28	44	79
			A2	****	****	****	71	75	86	87	96	99
JJA	****	****	A1B	****	****	****	70	75	90	82	93	98
			B1	****	****	****	52	63	72	61	76	87
			A2	****	****	****	50	67	81	73	87	96
SON	****	****	A1B	****	****	****	53	68	79	62	85	92
			B1	****	****	****	40	58	68	49	67	76
Frequency of Cold Days (TX10p)						<i>Future % frequency</i>						<i>Future % frequency</i>
Annual	****	****	A2	****	****	****	0	2	5	0	1	2
			A1B	****	****	****	0	3	4	0	1	3
			B1	****	****	****	1	4	6	1	3	4
			A2	****	****	****	0	2	5	0	0	1
DJF	****	****	A1B	****	****	****	0	2	4	0	0	3
			B1	****	****	****	0	3	6	0	3	3
			A2	****	****	****	0	2	4	0	0	2
MAM	****	****	A1B	****	****	****	0	2	3	0	0	2
			B1	****	****	****	0	4	6	0	3	3
			A2	****	****	****	0	0	2	0	0	0
JJA	****	****	A1B	****	****	****	0	0	1	0	0	1
			B1	****	****	****	0	0	1	0	0	3
			A2	****	****	****	0	1	2	0	0	1
SON	****	****	A1B	****	****	****	0	1	3	0	0	1
			B1	****	****	****	0	2	4	0	2	3
Frequency of Cold Nights (TN10p)						<i>Future % frequency</i>						<i>Future % frequency</i>
Annual	****	****	A2	****	****	****	1	3	5	0	0	2
			A1B	****	****	****	1	3	5	0	1	3
			B1	****	****	****	1	4	6	1	3	4
			A2	****	****	****	0	2	5	0	0	1
DJF	****	****	A1B	****	****	****	0	2	4	0	0	2
			B1	****	****	****	1	2	7	0	2	3
			A2	****	****	****	0	2	6	0	0	1
MAM	****	****	A1B	****	****	****	0	2	4	0	0	2
			B1	****	****	****	1	3	6	0	2	4
			A2	****	****	****	0	0	0	0	0	0
JJA	****	****	A1B	****	****	****	0	0	0	0	0	0
			B1	****	****	****	0	0	0	0	0	1
			A2	****	****	****	0	1	3	0	0	2
SON	****	****	A1B	****	****	****	0	1	4	0	0	2
			B1	****	****	****	1	3	5	0	2	4

	Observed Mean 1970-99	Observed Trend 1960-2006	Projected changes by the 2030s			Projected changes by the 2060s			Projected changes by the 2090s			
			Min	Median	Max	Min	Median	Max	Min	Median	Max	
			% total rainfall falling in Heavy Events (R95pct)									
	%	Change in % per decade					Change in %			Change in %		
Annual	****	****	A2	****	****	****	-5	0	5	-4	0	6
			A1B	****	****	****	-4	0	5	-4	0	4
			B1	****	****	****	-4	0	4	-3	1	4
DJF	****	****	A2	****	****	****	-5	0	12	-8	3	15
			A1B	****	****	****	-9	0	11	-6	1	12
			B1	****	****	****	-6	0	6	-7	2	9
MAM	****	****	A2	****	****	****	-15	-5	4	-18	-6	5
			A1B	****	****	****	-20	-4	6	-14	-6	2
			B1	****	****	****	-12	-3	5	-12	-2	5
JJA	****	****	A2	****	****	****	-8	-2	8	-19	-5	6
			A1B	****	****	****	-9	-3	8	-10	-4	6
			B1	****	****	****	-7	-1	5	-10	0	7
SON	****	****	A2	****	****	****	-6	1	7	-5	0	10
			A1B	****	****	****	-4	2	6	-5	-2	8
			B1	****	****	****	-4	0	6	-2	0	6
Maximum 1-day rainfall (RX1day)												
	mm	Change in mm per decade					Change in mm			Change in mm		
Annual	****	****	A2	****	****	****	-3	0	7	-3	1	8
			A1B	****	****	****	-4	0	9	-12	0	8
			B1	****	****	****	-4	0	7	-2	0	8
DJF	****	****	A2	****	****	****	-4	0	9	-4	2	10
			A1B	****	****	****	-6	0	12	-1	0	11
			B1	****	****	****	-3	-1	3	-2	1	6
MAM	****	****	A2	****	****	****	-5	-2	3	-7	-2	3
			A1B	****	****	****	-7	0	5	-10	-2	1
			B1	****	****	****	-6	-1	4	-8	0	2
JJA	****	****	A2	****	****	****	-5	-1	5	-6	-2	5
			A1B	****	****	****	-6	-1	2	-8	-1	2
			B1	****	****	****	-2	0	7	-8	-1	7
SON	****	****	A2	****	****	****	-2	0	9	-5	1	7
			A1B	****	****	****	-2	1	9	-8	0	6
			B1	****	****	****	-5	0	8	-2	0	10
Maximum 5-day Rainfall (RX5day)												
	mm	Change in mm per decade					Change in mm			Change in mm		
Annual	****	****	A2	****	****	****	-9	0	18	-9	2	25
			A1B	****	****	****	-7	0	19	-17	-3	16
			B1	****	****	****	-10	0	17	-5	2	26
DJF	****	****	A2	****	****	****	-7	-2	9	-11	1	17
			A1B	****	****	****	-13	0	16	-7	0	12
			B1	****	****	****	-8	-1	4	-4	2	13
MAM	****	****	A2	****	****	****	-14	-4	12	-21	-6	7
			A1B	****	****	****	-12	-5	6	-17	-6	4
			B1	****	****	****	-18	-1	5	-12	-2	6
JJA	****	****	A2	****	****	****	-9	-2	20	-19	-7	13
			A1B	****	****	****	-12	-4	9	-11	-5	13
			B1	****	****	****	-6	-2	14	-21	-2	18
SON	****	****	A2	****	****	****	-5	3	21	-16	3	14
			A1B	****	****	****	-4	5	21	-17	-4	13
			B1	****	****	****	-8	0	19	-8	2	19

* indicates trend is statistically significant at 95% confidence

**** indicates data are not available




Figure 1: Trends in annual and seasonal mean temperature for the recent past and projected future. All values shown are anomalies, relative to the 1970-1999 mean climate. Black curves show the mean of observed data from 1960 to 2006, Brown curves show the median (solid line) and range (shading) of model simulations of recent climate across an ensemble of 15 models. Coloured lines from 2006 onwards show the median (solid line) and range (shading) of the ensemble projections of climate under three emissions scenarios. Coloured bars on the right-hand side of the projections summarise the range of mean 2090-2100 climates simulated by the 15 models for each emissions scenario.




Figure 2: Spatial patterns of projected change in mean annual and seasonal temperature for 10-year periods in the future under the SRES A2 scenario. All values are anomalies relative to the mean climate of 1970-1999. In each grid box, the central value gives the ensemble median and the values in the upper and lower corners give the ensemble maximum and minimum.





Figure 3: Trends in monthly precipitation for the recent past and projected future. All values shown are anomalies, relative to the 1970-1999 mean climate. See Figure 1 for details.






Figure 5: Trends in monthly precipitation for the recent past and projected future. All values shown are percentage anomalies, relative to the 1970-1999 mean climate. See Figure 1 for details.




Figure 6: Spatial patterns of projected change in monthly precipitation for 10-year periods in the future under the SRES A2 scenario. All values are percentage anomalies relative to the mean climate of 1970-1999. See Figure 2 for details.




Figure 7: Trends in Hot-day frequency for the recent past and projected future. See Figure 1 for details.




Figure 8: Spatial patterns of projected change in Hot-day frequency for 10-year periods in the future under the SRES A2 scenario. See Figure 2 for details.




Figure 9: Trends in hot-night frequency for the recent past and projected future. See Figure 1 for details.




Figure 10: Spatial patterns of projected change in hot-night frequency for 10-year periods in the future under the SRES A2 scenario. See Figure 2 for details.




Figure 11: Trends in cold-day frequency for the recent past and projected future. See Figure 1 for details.




Figure 12: Spatial patterns of projected change in cold-day frequency for 10-year periods in the future under the SRES A2 scenario. See Figure 2 for details.




Figure 13: Trends in cold-night frequency for the recent past and projected future. See Figure 1 for details.




Figure 14: Spatial patterns of projected change in cold-night frequency for 10-year periods in the future under the SRES A2 scenario. See Figure 2 for details.




Figure 15: Trends in the proportion of precipitation falling in 'heavy' events for the recent past and projected future. All values shown are anomalies, relative to the 1970-1999 mean climate. See Figure 1 for details.




Figure 16: Spatial patterns of projected change in the proportion of precipitation falling in 'heavy' events for 10-year periods in the future under the SRES A2 scenario. All values are anomalies relative to the mean climate of 1970-1999. See Figure 2 for details.




Figure 17: Trends in maximum 1-day rainfall for the recent past and projected future. All values shown are anomalies, relative to the 1970-1999 mean climate. See Figure 1 for details.




Figure 18: Spatial patterns of maximum 1-day rainfall for 10-year periods in the future under the SRES A2 scenario. All values are anomalies relative to the mean climate of 1970–1999. See Figure 2 for details.




Figure 19: Trends in maximum 5-day rainfall for the recent past and projected future. All values shown are anomalies, relative to the 1970-1999 mean climate. See Figure 1 for details.




Figure 20: Spatial patterns of projected change in maximum 5-day rainfall for 10-year periods in the future under the SRES A2 scenario. All values are anomalies relative to the mean climate of 1970-1999. See Figure 2 for details.